

### Young pigeons

3-4 WEEKS INTERVAL

6 weeks of age	Rota-paramyxo
8 weeks of age	Salmonella/paratyphoid-herpes
10 weeks of age	<b>Smallpox</b> (injection/follicle method)
3-4 weeks before 1st training flights	<b>Rota-paramyxo BOOSTER</b>
Half June - early July*	Paramyxo-herpes <b>BOOSTER (HERPES)</b>

**TAILORED ADVICE**  
Contact us at  
[info@neornivet.be](mailto:info@neornivet.be)

**TIMING** The sequence and repetition of vaccines is partly determined by the infection pressure in your loft. NeorniVet will be happy to guide you through this.

**YOUNG PIGEON DISEASE** A **double vaccination** 3-4 weeks apart against **rotavirus** gives the best protection for young pigeon disease (often referred to as "adeno" in popular parlance). In case of heavy infection pressure, a **third booster** can be given 3 weeks before training. Vaccination against adenovirus and circovirus are not recommended, due to their unproven effectiveness and the lack of pigeon strains.

**SUPPORT** Support your young pigeons during the vaccination period in the first 6 weeks after weaning with **Immuno-Plus**, **Dysbac Liquid** and **Trichoryza Drops** from our product range **NeorniPharma**.



### Old pigeons & yearlings

Mid-February	Smallpox (injection/follicle method) <small>DISTINCT ADDED VALUE FOR YEARLINGS</small>
End of February	<b>Paramyxo-herpes</b>
14 days later	<b>Salmonella/paratyphoid**</b>

**SUPPORT** Guide your old pigeons and yearlings optimally to the start of the new season with **Trichoryza Drops**, **Dysbac Liquid** and **Ready** from our product range **NeorniPharma**.



### Breeders

6-7 weeks before breeding	<b>Salmonella/paratyphoid**</b>
Mid-October	<b>Paramyxo-herpes</b>

**SALMONELLA/PARATYPHOID** Depending on the infection pressure, it may or may not be necessary to treat the pigeons before vaccination and vaccinate once or twice.

**HERPES** In case of herpes problems during the rearing of young pigeons, it is recommended to vaccinate the breeding hens 2 times against herpes.

**PARAMYXO** An annual paramyxo vaccination is mandatory for breeders.

**SUPPORT** Stimulate the breeders 2 weeks before pairing up until the 1st clutch with **Vita-Min-DCP** and **Immuno-Plus** from our product range **NeorniPharma**. Bad fertilisers/old males: supplement with **Vit E+Se**.



\* 3-4 weeks before critical period - \*\* Optionally after preventive treatment in case of previous infection