



3-4 WEEKS INTERVAL

Young pigeons

6 weeks of age	Rota-paramyxo
8 weeks of age	Salmonella/paratyphoid-herpes
10 weeks of age	Smallpox (injection/follicle method)
3-4 weeks before 1st training flights	Rota-paramyxo BOOSTER
Half June - early July*	Paramyxo-herpes BOOSTER (HERPES)

TAILORED ADVICE
Contact us at
info@neornivet.be

TIMING The sequence and repetition of vaccines is partly determined by the infection pressure in your loft. NeornivEt will be happy to guide you through this.

YOUNG PIGEON DISEASE A **double vaccination** 3-4 weeks apart against **rotavirus** gives the best protection for young pigeon disease (often referred to as "adeno" in popular parlance). In case of heavy infection pressure, a **third booster** can be given 3 weeks before training. Vaccination against adenovirus and circovirus are not recommended, due to their unproven effectiveness and the lack of pigeon strains.

SUPPORT Support your young pigeons during the vaccination period in the first 6 weeks after weaning with **Immuno-Plus**, **Dysbac Liquid** and **Trichoryza Drops** from our product range **NeorniPharma**.



Old pigeons & yearlings

Mid-February	Smallpox (injection/follicle method) DISTINCT ADDED VALUE FOR YEARLINGS
End of February	Paramyxo-herpes
14 days later	Salmonella/paratyphoid**

SUPPORT Guide your old pigeons and yearlings optimally to the start of the new season with **Trichoryza Drops**, **Dysbac Liquid** and **Ready** from our product range **NeorniPharma**.



Breeders

6-7 weeks before breeding	Salmonella/paratyphoid**
Mid-October	Paramyxo-herpes

SALMONELLA/PARATYPHOID Depending on the infection pressure, it may or may not be necessary to treat the pigeons before vaccination and vaccinate once or twice.

HERPES In case of herpes problems during the rearing of young pigeons, it is recommended to vaccinate the breeding hens 2 times against herpes.

PARAMYXO An annual paramyxo vaccination is mandatory for breeders.

SUPPORT Stimulate the breeders 2 weeks before pairing up until the 1st clutch with **Vita-Min-DCP** and **Immuno-Plus** from our product range **NeorniPharma**. Bad fertilisers/old males: supplement with **Vit E+Se**.



* 3-4 weeks before critical period - ** Optionally after preventive treatment in case of previous infection